

References

- Ozek E, Soll R, Schimmel MS Partial exchange transfusion to prevent neurodevelopmental disability in infants with polycythemia. *Cochrane Database Syst Rev* 2010(1):CD005089.
- Mimouni FB, Merlob P, Dollberg S, et al. Neonatal polycythaemia: critical review and a consensus statement of the Israeli Neonatology Association. *Acta Paediatr* 2011;100(10):1290-6.
- Matthews DC, Glader B. Chapter 77 - Erythrocyte Disorders in Infancy A2 - Gleason, Christine A. In: S. U. Devaskar ed. *Avery's Diseases of the Newborn* (Ninth Edition). Philadelphia: W.B. Saunders; 2012:1080-107.
- Linderkamp, O., Versmold, H. T., Strohacker, I., Messow-Zahn, K., Riegel, K. P., & Betke, K. (1977). Capillary-venous hematocrit differences in newborn infants. I. Relationship to blood volume, peripheral blood flow, and acid base parameters. *Eur J Pediatr*, 127(1), 9-14. doi:10.1007/bf00465560
- Shohat, M., Merlob, P., & Reisner, S. H. (1984). Neonatal polycythemia: I. Early diagnosis and incidence relating to time of sampling. *Pediatrics*, 73(1), 7-10.
- Shohat, M., Reisner, S. H., Mimouni, F., & Merlob, P. (1984). Neonatal polycythemia: II. Definition related to time of sampling. *Pediatrics*, 73(1), 11-13.
- Drew, J. H., Guaran, R. L., Grauer, S., & Hobbs, J. B. (1991). Cord whole blood hyperviscosity: measurement, definition, incidence and clinical features. *J Paediatr Child Health*, 27(6), 363-365.
- Ramamurthy, R. S., & Berlanga, M. (1987). Postnatal alteration in hematocrit and viscosity in normal and polycythemic infants. *J Pediatr*, 110(6), 929-934.
- Bada, H. S., Korones, S. B., Pourcyrous, M., Wong, S. P., Wilson, W. M., 3rd, Kolni, H. W., & Ford, D. L. (1992). Asymptomatic syndrome of polycythemic hyperviscosity: effect of partial plasma exchange transfusion. *J Pediatr*, 120(4 Pt 1), 579-585. doi:10.1016/S0022-3476(05)82487-4
- Black, V. D., Lubchenco, L. O., Koops, B. L., Poland, R. L., & Powell, D. P. (1985). Neonatal hyperviscosity: randomized study of effect of partial plasma exchange transfusion on long-term outcome. *Pediatrics*, 75(6), 1048-1053.
- Goldberg, K., Wirth, F. H., Hathaway, W. E., Guggenheim, M. A., Murphy, J. R., Braithwaite, W. R., & Lubchenco, L. O. (1982). Neonatal hyperviscosity. II. Effect of partial plasma exchange transfusion. *Pediatrics*, 69(4), 419-425.
- Kumar, A., & Ramji, S. (2004). Effect of partial exchange transfusion in asymptomatic polycythemic LBW babies. *Indian Pediatr*, 41(4), 366-372.
- van der Elst, C. W., Molteno, C. D., Malan, A. F., & de V. Heese, H. (1980). The management of polycythaemia in the newborn infant. *Early Hum Dev*, 4(4), 393-403. doi:10.1016/0378-3782(80)90044-4
- UCSF Benioff Intensive Care Nursery Housestaff Manual: Polycythemia/Hyperviscosity. https://www.ucsfbenioffchildrens.org/pdf/manuals/39_Polycythemia.pdf