

References

1. AAP Clinical Practice Guideline. Urinary Tract Infection: clinical practice guideline for the diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. 2011. *Pediatrics*. 128,3: 595-610.
2. Marmor A. Evidence-based review of diagnosis and management of urinary tract infection in febrile infants/children <24 months of age. *UCSF Wiki*. March 2012.
3. National Institute for Health and Clinical Excellence. Urinary tract infection in children: NICE guideline. June 2010. <https://www.nice.org.uk/guidance/cg54>
4. The QuickWee trial: protocol for a randomised controlled trial of gentle suprapubic cutaneous stimulation to hasten non-invasive **urine collection** from **infants**. Kaufman J, Fitzpatrick P, Tosif S, Hopper SM, Bryant PA, Donath SM, Babl FE. *BMJ Open*. 2016 Aug 10;6(8):e011357. doi: 10.1136/bmjopen-2016-011357.

Published Children's Hospital Guidelines / Pathways

Children's Hospital of Philadelphia (CHOP)

Cincinnati Children's

Seattle Children's

UCSF BCH San Francisco, Pediatric Antimicrobial Stewardship Program BCH Community Acquired UTI Empiric Antibiotic Therapy Guidelines. <http://idmp.ucsf.edu/pediatric-guidelines-urinary-tract-infections-community-onset>